



King's Research Portal

DOI:

[10.1371/journal.pone.0034821](https://doi.org/10.1371/journal.pone.0034821)

Document Version

Publisher's PDF, also known as Version of record

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Prina, M., Deeg, D., Brayne, C., Beekman, A., & Huisman, M. (2012). The Association between Depressive Symptoms and Non-Psychiatric Hospitalisation in Older Adults. *PLoS One*, 7(4), N/A. [e34821].
<https://doi.org/10.1371/journal.pone.0034821>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

The Association between Depressive Symptoms and Non-Psychiatric Hospitalisation in Older Adults

A. Matthew Prina^{1*}, Dorly Deeg^{2,3}, Carol Brayne¹, Aartjan Beekman^{2,3}, Martijn Huisman^{2,3,4}

1 Department of Public Health & Primary Care, Institute of Public Health, University of Cambridge, Cambridge, United Kingdom, **2** Department of Epidemiology & Biostatistics and the EMGO Institute for Health and Care Research, VU University Medical Center, Amsterdam, The Netherlands, **3** Department of Psychiatry, VU University Medical Center, Amsterdam, The Netherlands, **4** Department of Sociology, VU University, Amsterdam, The Netherlands

Abstract

Background: It is known that people who suffer from depression are more likely to have other physical illnesses, but the extent of the association between depression and non-psychiatric hospitalisation episodes has never been researched in great depth. We therefore aimed to investigate whether depressed middle-aged and older people were more likely to be hospitalised for causes other than mental illnesses, and whether the outcomes for this group of people were less favourable.

Methods & Findings: Hospital events from 1995 to 2006 were obtained from the Dutch National Medical Register and linked to participants of the Longitudinal Aging Study Amsterdam (LASA). Linkage was accomplished in 97% of the LASA sample by matching gender, year of birth and postal code. Depression was measured at each wave point of the LASA study using the Centre for Epidemiologic Studies Depression (CES-D). Hospital outcomes including admission, length of stay, readmission and death while in hospital were recorded at 6, 12 and 24 months intervals after each LASA interview. Generalised Estimating Equation models were also used to investigate potential confounders. After 12 months, 14% of depressed people were hospitalised compared to 10% of non-depressed individuals. There was a 2-fold increase in deaths while in hospital amongst the depressed (0.8% vs 0.4%), who also had longer total length of stay (2.6 days vs 1.4 days). Chronic illnesses and functional limitations had major attenuating effects, but depression was found to be an independent risk factor for length of stay after full adjustment (OR = 1.33, 95% CI: 1.22–1.46 after 12 months).

Conclusions: Depression in middle and old age is associated with non-psychiatric hospitalisation, longer length of stay and higher mortality in clinical settings. Targeting of this high-risk group could reduce the financial, medical and social burden related to hospital admission.

Citation: Prina AM, Deeg D, Brayne C, Beekman A, Huisman M (2012) The Association between Depressive Symptoms and Non-Psychiatric Hospitalisation in Older Adults. PLoS ONE 7(4): e34821. doi:10.1371/journal.pone.0034821

Editor: Jerson Laks, Federal University of Rio de Janeiro, Brazil

Received: February 4, 2012; **Accepted:** March 5, 2012; **Published:** April 4, 2012

Copyright: © 2012 Prina et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The LASA study was primarily funded by the Ministry of Health, Welfare and Sports (to The Longitudinal Aging Study Amsterdam). The funders had no role in the study design, data collection, analysis, decision to publish or preparation of the manuscript. No additional external funding was received for this study.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: amp68@medschl.cam.ac.uk

Introduction

Although adults aged 65 and over encompass for only a small percentage of global populations, they account for over 45% percent of all hospital admissions in England [1]. A rise in the number of older adults will have dramatic consequences on the already strained and saturated national health services. For that reason, maintaining individuals in home settings, reducing preventable admissions, has been on the agendas of UK health policy makers for several years, yet population evidence concerning major contributors of hospitalisations is still not well understood.

One of the largest contributors to the worldwide burden of morbidity is depression [2], which might be related to a considerable part of hospital admissions in developed countries too. Depression is a common mental disorder that is widespread amongst the older population and that is associated with physical co-morbidities, lower quality of life and disability [3,4,5]. Although it is known that there is a strong association between depression and chronic conditions, it is not yet clear whether this indeed

translates into higher numbers of hospital admissions and worse hospital outcomes, as no large epidemiological studies have been conducted to investigate this.

The absence of evidence can partially be blamed on lack of comprehensive medical registers that are linkable to other datasets. Medical registers on their own have in fact poor sensitivity for depression, as a large majority of depressed people living in the community are not formally diagnosed and their mental health statuses are not recorded. Therefore, an external population representative dataset, where mental health status is measured, is essential for this type of research. The existence of a national medical register in the Netherlands, where data linkage was possible, allowed us to carry out this research.

Several studies have investigated the relationship between depression and hospital outcomes in specific populations (e.g. cardiovascular patients, diabetes patients, etc.) [6,7,8,9,10] and most have reported positive associations, however depression was measured at admission in the majority of these. To our knowledge, this is the only study that measured depressive symptoms prospectively without focusing specifically on one cause of hospital




Figure 1. Number of linkable LASA records by wave and cohort.
doi:10.1371/journal.pone.0034821.g001

admission, and following a large number of middle-aged and older participants for up to 24 months.

We aimed to assess whether depressed middle-aged and older adults were more likely to be hospitalised for causes other than mental illnesses. We also assessed differences in hospital outcomes, namely length of stay, hospital mortality and re-admissions between depressed and non-depressed individuals.

Methods

Sources of data

Participants were selected from the Longitudinal Aging Study Amsterdam (LASA) [11], a longitudinal study that includes a representative random sample of Dutch adults aged between 55 and 85 at baseline. The methods and characteristics of this sample have been described in great detail in previous publications [12]. Data collection in LASA started in 1992–1993 (LASA baseline) with follow-ups at roughly 3 years intervals. In 2003/2003, ten years after the baseline measurement of the first LASA cohort, a new cohort was started with adults aged between 55 and 65 years at that time. This new population sample was selected using the same procedures and sampling frame as the initial LASA cohort. Respondents of both the initial and the second LASA cohort have been followed-up together since then.

To address our research question we included LASA participants who had at least one interview between 1995 and 2006, because access to complete medical records was only available for that time frame. A total of 8560 observations across different waves and cohorts were recorded, as illustrated in **figure 1**.

The Dutch Hospital Discharge Register (HDR) [13] is a register of hospital admissions. All general and academic hospitals and most single-specialty hospitals, with the exception of epilepsy clinics and long-stay centres for rehabilitation and asthma treatment, are included in this register. Thus, the register provides a nearly complete coverage of hospital inpatient treatments in the Netherlands. All clinical and day admissions are registered based on a uniform registration system. The data include admission and discharge dates, extensive treatment information (including about 10,000 medical procedures), and death.

Information from the HDR about individuals can be linked to other datasets via the Municipal Population Registration (MPR). The MPR contains basic information like date of birth, registered

partners, and postal code for everyone listed in a Dutch municipality. Linkage of the HDR to the MPR occurred on the basis of a set of identifying variables: postal code, date of birth and gender [13]. The LASA dataset was also linked to the MPR, by Statistics Netherlands especially for our research, so that LASA data could be linked with information from the HDR as well. Linkage of LASA data with the MPR also occurred on the basis of identifying background variables. Participants from LASA were linked to data from the HDR at each wave and hospitalisation was followed up for 6, 12 and 24 months. Only records that had a full 24 months follow up, or censored data due to death, were included. Linkage was fully achieved for 8282 records, 97% of the initial sample. Consent forms for accessing medical records were signed by the LASA participants and the study was approved by the Medical Ethical Committee of the VU University Medical Centre.

Measures and outcomes

Depression, physical co-morbidities, socio-demographics and functional limitations were recorded in LASA during each wave.

Depressive symptoms were measured at each wave using the Centre for Epidemiological Studies Depression Scale (CES-D), using a cut-off score of 16 points. This scale has been validated and been found to have good specificity and sensitivity for depression and to be a satisfactory instrument to measure depression severity in older people [14,15,16], with similar psychometric properties in its Dutch version [17]. Criterion validity for major depression was also shown to be very high, with a sensitivity of 100% and a specificity of 88% [18], and with very little overlap with symptoms of physical morbidity [19,20].

Socio-demographics co-variables included *age*, *sex* and *level of education*. This was classified as no education, elementary/lower, intermediate or higher education. The total number of *chronic illnesses* ranged from 0 to 9 and was identified by asking questions on arthritis, peripheral atherosclerosis, cardiac disease, diabetes, malignant neoplasms, chronic lung disease, stroke, and a maximum of two other chronic diseases. These self-reported answers of LASA respondents were shown to correspond well with information from general practitioners [21]. A questionnaire on difficulty experienced with several activities (e.g. cutting own toenails, using public transport and walking up & down stairs) was used to

measure *functional limitations*. These three questions were selected from the validated OECD Questionnaire [22], after showing good internal reliability (0.73) and being a good proxy for functional limitations [23]. Functional limitations were categorised into four severity groups according to the response to the three questions.

Smoking was categorised in current use, past use or no use. Alcohol consumption was measured with a questionnaire developed for the Netherlands Health Interview Survey [24] and was classified into two categories (excessive/severe vs not excessive/severe) according to the Garretsen Index of Present Alcohol Use [25]. Excessive alcohol use is defined as drinking at least once a week six glasses or more, or 21 days per month drinking four or more glasses.

The *outcomes* that were investigated were hospital admission (0 = non admitted, 1 = admitted into hospital), mean total length of stay across the different hospitalisation episodes, death while in hospital, number of readmissions (excluding no admissions) and type of admission. This latter was either day care admission (i.e. day hospital) or full overnight admission.

Statistical analyses

After pooling the data from each wave and cohort, we tabulated hospital outcomes by depression status for three follow-up periods: 6, 12 and 24 months. Differences in the outcome variables by depression status were evaluated with Chi-Square χ^2 tests for categorical variables.

Due to the longitudinal nature of our observations, Generalised Estimating Equations (GEE), which take into account the dependency of the repeated measurements within persons, were used to investigate risk factors. Based on the nature and distribution of the outcome we decided what type of GEE regression function to use (e.g. logit vs negative binomial).

All the models were initially adjusted for age and sex, before fully adjusted models were run. The most conservative models are reported. In order to determine whether a potential confounder should be included in the final model, we added one factor at the time to see whether it showed a significant association with one or more of the outcomes.

Moreover, we selected all the participants ($n = 2251$) who had fully linkable hospital events from the first LASA cohort (wave 1995–1996) and plotted Kaplan-Meier survival curves (36 months) using hospital admission and death in hospital as outcomes. Data was only gathered from the first cohort to maintain a high consistency and to exclude that introduction of health policies or changes in health services could affect the results. Moreover a full follow up of 36 months was not present for all participants in the second cohort. Log rank tests were also performed to compare the cumulative admission and mortality rates between depressed and non-depressed individuals.

Results

Depressed older adults were more likely to be older, be women, have lower socio-economic status, and have poorer physical health than individuals free of depressive symptoms (**table 1**). Two years after the interview, 27% of depressed older adults were admitted into hospital at least once, compared to only 23% of non-depressed participants (**table 2**). These differences were statistically significant also at 12 and 6 months, as confirmed by the χ^2 p values, which were lower than 0.0010. Depressed older adults were also twice more likely to die while in hospital than non-depressed individuals (p values < 0.05).

There were not any major differences in the number of readmissions at any of the follow-up times. However, there were

Table 1. Population characteristics of linkable observations.

Characteristics	No Depressive symptoms (frequency)		Depressive symptoms (frequency)	
Age group				
54–65	2161	(30.99)	314	(24.01)
65–74	2287	(32.79)	336	(25.69)
74–85	1925	(27.60)	452	(34.56)
85+	601	(8.62)	206	(15.75)
Sex				
Male	3383	(48.51)	391	(29.89)
Female	3591	(51.49)	917	(70.11)
Education				
No	2189	(31.39)	609	(46.67)
Elementary/Lower	3486	(49.99)	539	(41.30)
Intermediate	962	(13.80)	125	(9.58)
Higher	336	(4.82)	32	(2.45)
Smoking				
Never	1786	(31.71)	400	(36.53)
Used to	2820	(50.07)	453	(41.37)
Smokes	1026	(18.22)	242	(22.10)
Alcohol (potential alcohol problem)				
No	5366	(95.28)	1052	(96.07)
Yes	266	(4.72)	43	(3.93)
Functional Limitations				
No	3755	(54.79)	288	(22.41)
2 without	1572	(22.82)	249	(19.38)
1 without	947	(13.74)	321	(24.98)
All without	596	(8.65)	427	(33.23)
Chronic illnesses				
Mean number	1.67	(1.64–1.70)	2.50	(2.41–2.58)
Anxiety				
No	5831	(95.14)	555	(47.76)
Yes	298	(4.86)	607	(52.54)

doi:10.1371/journal.pone.0034821.t001

marked variations in mean total length of stay between the two groups. At 24 months, hospitalised depressed individuals spent on average 5.3 (95% CI: 4.4–6.3) days in hospital compared to only 3.5 days (95% CI: 3.2–3.8) for participants without depression. Similar differences were found at 6 and 12 months (**table 2**). The last outcome that was investigated was related to the type of admission. Two types of admission were recorded: day hospital or full admission. Differences in the types of admissions were found at 6 months (p value < 0.05) but not at 12 ($p = 0.07$) or 24 months ($p = 0.09$). Although during the latter two follow-up periods the variations were only marginally significant depressed older adults had a lower ratio of day hospital to full admissions compared to their depressed peers.

Confounders and associated risk factors

Generalised Estimating Equations [26,27] were used to investigate potential risk factors involved in explaining the association between depression and hospital outcomes (**table 3**).

Table 2. Hospital admissions and other hospital outcomes for depressed (CES-D>16) and non-depressed (CES-D<16) participants.

	24 Months				12 Months				6 Months			
Hospital outcomes	Not Depressed		Depressed		Not Depressed		Depressed		Not Depressed		Depressed	
Hospital Admission												
No (frequency)	5371	(77.01)	952	(72.78)	6252	(89.65)	1131	(86.47)	6461	(92.64)	1171	(89.53)
Yes (frequency)	1603	(22.99)	356	(27.22)	722	(10.35)	177	(13.53)	513	(7.36)	137	(10.47)
Death in hospital												
No (frequency)	6866	(98.45)	1270	(97.09)	6946	(99.6)	1297	(99.16)	6957	(99.76)	1299	(99.31)
Yes (frequency)	108	(1.55)	38	(2.91)	28	(0.40)	11	(0.84)	17	(0.24)	9	(0.69)
Day Care												
No (frequency)	1179	(73.55)	278	(78.09)	507	(70.22)	137	(77.40)	346	(67.45)	105	(76.64)
Yes (frequency)	424	(26.45)	78	(21.91)	215	(29.78)	40	(22.60)	167	(32.55)	32	(23.36)
Number of readmissions												
Mean (95% CI)	2.10	(1.99–2.20)	2.25	(2.02–2.48)	1.96	(1.84–2.08)	1.99	(1.77–2.22)	1.86	(1.74–1.98)	1.90	(1.66–2.13)
Total length of stay												
Mean (95% CI)	3.51	(3.21–3.81)	5.31	(4.37–6.26)	1.40	(1.21–1.59)	2.62	(1.98–3.25)	0.90	(0.76–1.05)	1.73	(1.23–2.23)

doi:10.1371/journal.pone.0034821.t002

Unadjusted Odds Ratios (ORs) for the depressed group were highest for the mortality outcome (6 months ORs = 2.8, 95% CI: 1.26–6.37), but were also above one for all the other outcomes. After adjustment for age and sex, depression was still found to be an independent risk factor for hospital admission, longer length of stay and death. After full adjustment for socio-economic status and physical co-morbid conditions, the effect of depression was attenuated. However, the ORs remained high for the mortality outcome, and depression was still found to be an independent risk factor for total length of stay (6 months ORs = 1.17, 95% CI:

1.07–12.9; 12 months ORs = 1.33, 95% CI: 1.22–1.46). Similar outcomes were found at 24 months follow up (**data not shown**). Separate regressions were also fitted to assess what factor led to the largest attenuation of the associations. Adjusting for the number of functional limitations consistently had the biggest impact on the Odds Ratios (**data not shown**).

Hospital admissions and deaths while in hospital were significantly different between groups on Kaplan-Meier analyses (**figure 2**). The p values of log-rank tests for the hospital admission and death were respectively equal to 0.002 and 0.008.

Table 3. GEE (Generalised Estimating Equation) models for the independent effect of depression (CES-D>16) on different outcomes.

Outcomes	Hospital Admission (function type: logistic)				Total Length of Stay (function type: negative binomial)				Dying During Hospital Stay (function type: negative binomial)			
	12 months		6 months		12 months		6 months		12 months		6 months	
Model 1												
Depression	1.35	(1.13–1.62)	1.47	(1.20–1.80)	1.86	(1.73–2.00)	1.88	(1.74–2.03)	2.10	(1.05–4.24)	2.84	(1.26–6.37)
Model 2												
Depression	1.34	(1.11–1.60)	1.47	(1.20–1.81)	1.79	(1.67–1.93)	1.78	(1.65–1.93)	1.96	(0.96–4.02)	2.67	(1.16–6.15)
Age	1.02	(1.01–1.03)	1.02	(1.01–1.03)	1.04	(1.04–1.04)	1.04	(1.04–1.04)	1.05	(1.02–1.09)	1.08	(1.03–1.13)
Sex	0.78	(0.67–0.90)	0.77	(0.65–0.91)	0.87	(0.82–0.92)	0.87	(0.81–0.92)	0.62	(0.33–1.18)	0.42	(0.18–0.94)
Model 3												
Depression	1.01	(0.83–1.22)	1.07	(0.86–1.33)	1.33	(1.22–1.46)	1.17	(1.07–1.29)	1.35	(0.58–3.17)	1.90	(0.68–5.28)
Age	1.00	(1.00–1.01)	1.00	(0.99–1.01)	1.04	(1.03–1.04)	1.04	(1.04–1.05)	1.20	(0.97–1.06)	1.04	(0.98–1.11)
Sex	0.70	(0.60–0.82)	0.69	(0.58–0.82)	0.80	(0.74–0.86)	0.80	(0.74–0.87)	0.70	(0.32–1.53)	0.58	(0.21–1.56)
Education	1.00	(0.91–1.10)	0.98	(0.88–1.10)	0.93	(0.89–0.98)	0.94	(0.89–0.98)	0.87	(0.53–1.44)	0.80	(0.42–1.53)
Chronic illnesses	1.20	(1.14–1.27)	1.23	(1.16–1.31)	1.16	(1.13–1.20)	1.23	(1.19–1.26)	1.21	(0.93–1.57)	1.18	(0.85–1.65)
Functional limitations	1.25	(1.15–1.35)	1.26	(1.14–1.38)	1.29	(1.24–1.33)	1.30	(1.24–1.35)	1.46	(0.98–2.18)	1.42	(0.86–2.38)
Smoking	n/a		n/a		1.20	(1.14–1.26)	1.20	(1.14–1.28)	1.93	(1.12–3.32)	2.13	(1.07–4.22)
Alcohol problem	n/a		n/a		1.29	(1.10–1.52)	1.43	(1.20–1.70)	0.70	(0.92–5.41)	1.28	(0.16–10.36)

doi:10.1371/journal.pone.0034821.t003




Figure 2. Kaplan-Meier survival curves for hospitalisation and death while in hospital. Red line: depressive symptoms (CES-D > 16). Blue line: no depressive symptoms (CES-D < 16). doi:10.1371/journal.pone.0034821.g002

Discussion

This is the first study of this scale that clearly identifies an association between depressive symptoms and non-psychiatric hospitalisation episodes and that does not specifically focus on specific hospitals, wards or diseases.

Our study confirmed that depression is associated with hospital admission and with worse hospital outcomes in a middle-aged and older depressed Dutch population. By introducing covariates in our models, the Odds Ratios were reduced between 45–97%. The association between depression and admission was mostly explained by higher physical co-morbidities and functional limitations but depression remained an independent risk factor for longer length of stay, and had a strong association with mortality as well. Several explanations could be put forward to explain the association between depression and poorer hospital outcomes.

It is known that depressed individuals often do not follow medical recommendations for underlying medical illnesses and have lower treatment compliance [28,29,30]. Although this is likely not to be of major concern during hospitalisation, it could affect those individuals while in the community. Those persons may consequently present to a hospital with a later or more severe stage of disease that requires more invasive treatment and a higher chance of adverse outcome that would affect length of stay and mortality. Unfortunately we could not tease out severity or type of treatment received during hospitalisation, but the fact that more depressed participants were admitted for full overnight admissions rather than for day care seems to suggest that acuteness and severity might be higher in this group. It is also possible that longer LOS may be the result of the discovery of the mental disorder during admission, requiring the involvement of liaison psychiatry services that could lengthen the stay, while trying to optimise the psychiatric illness.

The most apparent explanation for this association lies in the strength of association between depression and chronic conditions. Several mediators (e.g. limbic-hypothalamic-pituitary-adrenal axis alterations, pro-inflammatory milieu, decline in glucocorticoid receptor function, etc.) occurring in depressed individuals that increase the likelihood of physical morbidity have been described by Wolkowitz OM, et al. [31]. However, our findings suggest that the relationship between depression and hospitalisation is not solely mediated by chronic conditions and might result from the synergic action of several different pathways. Our findings also

suggest that functional limitations may be on the pathway to hospital outcomes in depressed patients, or conversely, that functional limitations are causing depressive symptoms that in turn lead to poor hospital outcomes. The causal association between depression and disability has been shown to be bidirectional [32,33].

Strengths

The major strengths of our study stand in the large cohort, in its longitudinal nature, and in the excellent validity of the linked registry. The latter was demonstrated in a random sample of records, showing that 84% of the main diagnoses (validated through medical record review by medical specialists) and 99% of the personal, admission and discharge data were correctly recorded in the hospital discharge register [34]. The longitudinal nature of the study allowed us to identify depression first, and assess hospital outcomes thereafter, thus avoiding ambiguities in causal direction. No studies have been able to investigate the association between depression and hospital outcomes in such detail and with such complete coverage.

Limitations

This is a study conducted in a nationally representative sample of the Dutch middle-aged and older population. We believe that our findings could be easily applicable to other western developed countries. Even though there are differences in the prevalence of lifetime depression and impairment related to depression between western developed countries [35], the burden of morbidity due to depression is immense in all of them. However specifics about access to health care, treatment, and recognition of depression in older adults are likely to differ [36], which may affect the likelihood of depressed older adults to become hospitalised.

Depressive symptoms were only measured at the beginning of each wave. Considering the episodic nature of this disorder and the length of our follow up (up to 24 months), we are therefore not able to exclude classification bias (e.g. some participants free from the condition could have developed depression before hospitalisation, whereas remission could have occurred amongst some depressed participants).

The CES-D is a symptom rating scale, and although this measures clinically relevant depressive symptoms, it is not comparable to the definition used in the DSM classification

[37]. Nevertheless, it has been demonstrated that the consequences of depressive symptoms in older adults are very similar to those of major depressive illnesses [38].

Finally, using a general population sample, compared to a sample with specific medical conditions, has its advantages but we should also emphasise its limitations. In a more homogenous sample a comparison of those depressed and not depressed is less likely to be confounded by differences in co-occurring medical conditions. Moreover, in this highly selected groups, measuring and controlling for the severity of the medical illness is easier to make compared to a general cohort.

Implications & Conclusions

Although depression is so prevalent in middle-aged and older adults and has adverse effects on hospital outcomes, it is not often noticed and treated in general hospitals. Clinicians should be aware of the high prevalence of depression in older adults, and also of the importance of treatment options and early identification of depressive symptoms that could reduce hospitalisation, length of stay and improve survival.

It is known that the bulk of hospitalisations occur among older adults and that hospital care is the most costly form of care. Our findings strongly suggest that there are opportunities to drastically reduce costs related to admissions if case finding and treatment of depressive symptoms in older adults could be improved. Further evidence to support the previous claim is certainly needed as

although it is known that screening paired with an organised system of depression care can improve depression outcomes [39], it is not clear whether physical outcomes could also be improved. Future research that includes a full economic analysis of this prevention approach is needed, together with studies aimed at investigating the role of clinical depression but also of anxiety, so often co-morbid with depression. A trial for case finding and effective treatment of depression in a high-risk group of older adults (e.g. comorbid depression and chronic medical illnesses), to see whether hospital outcomes could be improved, is also warranted. Although we solely focused on one direction of this association, this relationship is likely to be bi-directional, where those sick enough to be hospitalised are more often depressed, potentially creating a feedback loop between mental status and admissions into hospital. The investigation of this phenomenon deserves more attention. In conclusions, this study suggests that there is an association between depression, hospital admission and worse hospital outcomes, identifying a potential target to reduce preventable admissions and ensuing associated costs.

Author Contributions

Conceived and designed the experiments: AMP DD CB AB MH. Performed the experiments: AMP MH. Analyzed the data: AMP MH. Contributed reagents/materials/analysis tools: CB DD AB MH. Wrote the paper: AMP MH CB AB.

References

- Department of Health (2010–2011) Hospital Episode Statistics (HES).
- Lopez AD, Mathers CD, Ezzati M, Jamison DT, Murray CJ (2006) Global and regional burden of disease and risk factors, 2001: systematic analysis of population health data. *Lancet* 367: 1747–1757.
- Knol MJ, Twisk JW, Beekman AT, Heine RJ, Snoek FJ, et al. (2006) Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis. *Diabetologia* 49: 837–845.
- Wulsin LR, Singal BM (2003) Do depressive symptoms increase the risk for the onset of coronary disease? a systematic quantitative review. *Psychosomatic Medicine* 65: 201–210.
- Covinsky KE, Yaffe K, Lindquist K, Cherkasova E, Yelin E, et al. (2010) Depressive symptoms in middle age and the development of later-life functional limitations: the long-term effect of depressive symptoms. *J Am Geriatr Soc* 58: 551–556.
- Turner A, Phillips L, Hambridge JA, Baker AL, Bowman J, et al. (2010) Clinical outcomes associated with depression, anxiety and social support among cardiac rehabilitation attendees. *Aust N Z J Psychiatry* 44: 658–666.
- Kato N, Kinugawa K, Yao A, Hatano M, Shiga T, et al. (2009) Relationship of depressive symptoms with hospitalization and death in Japanese patients with heart failure. *J Card Fail* 15: 912–919.
- Eisner MD, Katz PP, Lactao G, Iribarren C (2005) Impact of depressive symptoms on adult asthma outcomes. *Annals of Allergy, Asthma & Immunology* 94: 566–574.
- Hosaka T, Aoki T, Watanabe T, Okuyama T, Kurosawa H (1999) Comorbidity of depression among physically ill patients and its effect on the length of hospital stay. *Psychiatry and Clinical Neuroscience* 53: 491–495.
- Frasure-Smith N, Lesperance F, Gravel G, Masson A, Juneau M, et al. (2000) Depression and health-care costs during the first year following myocardial infarction. *J Psychosom Res* 48: 471–478.
- Deeg D, Knipscheer C, Van Tilburg W (2003) Autonomy and Well-being in the Aging Population: Concepts and Design of the Longitudinal Aging Study Amsterdam. In: Netherlands Institute of Gerontology, editor (2003) Bunnik, The Netherlands.
- Huisman M, Poppelaars J, van der Horst M, Beekman AT, Brug J, et al. (2011) Cohort Profile: The Longitudinal Aging Study Amsterdam. *Int J Epidemiol*.
- de Bruin A, Kardaun J, Gast F, de Bruin E, van Sijl M, et al. (2004) Record linkage of hospital discharge register with population register: Experiences at Statistics Netherlands. *Statistical Journal of the United Nations* 21: 23–32.
- Himmelfarb S, Murrell SA (1983) Reliability and validity of five mental health scales in older persons. *J Gerontol* 38: 333–339.
- Radloff LS, Teri L (1986) Use of the CES-D with older adults. *Clinical Gerontology* 5: 119–136.
- Hertzog C VAJ, Usala PD, Hultsch DF, Dixon R (1990) Measurement properties of the Center for Epidemiological Studies Depression Scale (CES-D) in older populations. *Psychological Assessment* 2: 64–72.
- Beekman AT, van Limbeek J, Deeg DJ, Wouters L, van Tilburg W (1994) [A screening tool for depression in the elderly in the general population: the usefulness of Center for Epidemiological Studies Depression Scale (CES-D)]. *Tijdschr Gerontol Geriatr* 25: 95–103.
- Beekman AT, Deeg DJ, Van Limbeek J, Braam AW, De Vries MZ, et al. (1997) Criterion validity of the Center for Epidemiologic Studies Depression scale (CES-D): results from a community-based sample of older subjects in The Netherlands. *Psychological Medicine* 27: 231–235.
- Berkman LF, Berkman CS, Kasl S, Freeman DH, Jr., Leo L, et al. (1986) Depressive symptoms in relation to physical health and functioning in the elderly. *Am J Epidemiol* 124: 372–388.
- Foelker GA, Jr., Shewchuk RM (1992) Somatic complaints and the CES-D. *J Am Geriatr Soc* 40: 259–262.
- Kriegsman DM, Penninx BW, van Eijk JT, Boeke AJ, Deeg DJ (1996) Self-reports and general practitioner information on the presence of chronic diseases in community dwelling elderly. A study on the accuracy of patients' self-reports and on determinants of inaccuracy. *J Clin Epidemiol* 49: 1407–1417.
- Van Sonsbeek J (1988) Methodological and substantial aspects of the OECD indicator of chronic functional limitations. *Maandbericht Gezondheid (CBS)* 88: 4–17.
- Smits CH, Deeg DJ, Jonker C (1997) Cognitive and emotional predictors of disablement in older adults. *J Aging Health* 9: 204–221.
- (1993) Statistical Yearbook. In: (CBS) CBoS, editor (1993) Den Haag.
- Garretsen H, ed. Probleemdrinken: Prevalentiebepaling, beïnvloedende factoren en preventiemogelijkheden: Theoretische overwegingen en onderzoek in Rotterdam. Lisse.
- Liang K, Zeger S (1993) Regression analysis for correlated data. *Annual Reviews of Public Health* 14: 43–68.
- Twisk JW (1997) Different statistical models to analyze epidemiological observational longitudinal data: an example from the Amsterdam Growth and Health Study. *Int J Sports Med* 18 Suppl 3: S216–224.
- Ziegelstein RC, Fauerbach JA, Stevens SS, Romanelli J, Richter DP, et al. (2000) Patients with depression are less likely to follow recommendations to reduce cardiac risk during recovery from a myocardial infarction. *Arch Intern Med* 160: 1818–1823.
- Kronish IM, Rieckmann N, Halm EA, Shimbo D, Vorchheimer D, et al. (2006) Persistent depression affects adherence to secondary prevention behaviors after acute coronary syndromes. *J Gen Intern Med* 21: 1178–1183.
- Katon WJ (2011) Epidemiology and treatment of depression in patients with chronic medical illness. *Dialogues Clin Neurosci* 13: 7–23.
- Wolkowitz OM, Reus VI, Mellon SH (2011) Of sound mind and body: depression, disease, and accelerated aging. *Dialogues Clin Neurosci* 13: 25–39.
- Ormel J, Vonkorff M, Oldehinkel AJ, Simon G, Tiemens BG, et al. (1999) Onset of disability in depressed and non-depressed primary care patients. *Psychological Medicine* 29: 847–853.

33. Penninx BW, Deeg DJ, van Eijk JT, Beekman AT, Guralnik JM (2000) Changes in depression and physical decline in older adults: a longitudinal perspective. *Journal of Affective Disorders* 61: 1–12.
34. Paas G, Veenhuizen K (2002) Research on the validity of the LMR. Utrecht: Prismant.
35. Bromet E, Andrade LH, Hwang I, Sampson NA, Alonso J, et al. (2011) Cross-national epidemiology of DSM-IV major depressive episode. *BMC Med* 9: 90.
36. Braam AW, Prince MJ, Beekman AT, Delespaul P, Dewey ME, et al. (2005) Physical health and depressive symptoms in older Europeans. Results from EURODEP. *The British journal of psychiatry : the journal of mental science* 187: 35–42.
37. American Psychiatric Association (1994) *Diagnostic and Statistical Manual of Mental Disorders*. Fourth edition. Washington, DC: American Psychiatric Association.
38. Beekman AT, Deeg DJ, Braam AW, Smit JH, Van Tilburg W (1997) Consequences of major and minor depression in later life: a study of disability, well-being and service utilization. *Psychol Med* 27: 1397–1409.
39. Seedat S, Scott KM, Angermeyer MC, Berglund P, Bromet EJ, et al. (2009) Cross-national associations between gender and mental disorders in the World Health Organization World Mental Health Surveys. *Arch Gen Psychiatry* 66: 785–795.